

## THE STANDARD CANTOR FUNCTION IS SUBADDITIVE

JOZEF DOBOŠ

(Communicated by Andreas R. Blass)

ABSTRACT. In this paper the subadditivity of the Cantor function  $\phi: [0, 1] \rightarrow [0, 1]$  is proved.

Let us begin by recalling that a function  $f: \mathbb{R} \rightarrow \mathbb{R}$  is said to be subadditive, if it satisfies the inequality  $f(x + y) \leq f(x) + f(y)$  whenever  $x, y \in \mathbb{R}$ . (See [2] and [4].)

The usual definition of the standard Cantor function involves the classic middle-thirds description of the standard Cantor set. (See [1] and [3].) We offer an alternate definition of this function.

Define a sequence of functions  $\phi_n: \mathbb{R} \rightarrow [0, 1]$  by

$$\phi_0(x) = \begin{cases} 0 & \text{if } x \leq 0, \\ x & \text{if } 0 \leq x \leq 1, \\ 1 & \text{if } x \geq 1, \end{cases} \quad \phi_{n+1}(x) = \begin{cases} \frac{1}{2} \cdot \phi_n(3x) & \text{if } x \leq \frac{2}{3}, \\ \frac{1}{2} + \frac{1}{2} \cdot \phi_n(3x - 2) & \text{if } x \geq \frac{1}{3}. \end{cases}$$

It is easy to check that each  $\phi_n$  is non-decreasing, that  $\phi_n(x) = 0$  for all  $x \leq 0$ , that  $\phi_n(x) = 1$  for all  $x \geq 1$ , and that the two lines in the definition of  $\phi_{n+1}$  agree in the overlap of their domains, both giving  $\phi_{n+1}(x) = \frac{1}{2}$  when  $\frac{1}{3} \leq x \leq \frac{2}{3}$ .

Put  $\phi = \lim_{n \rightarrow +\infty} \phi_n$ . It is not difficult to verify that the restriction of  $\phi$  to  $[0, 1]$  is the standard Cantor function. The functions  $\phi_n$  are polygonal approximations of  $\phi$ .

**Theorem.** *The standard Cantor function is subadditive.*

*Proof.* The function  $\phi$  is the pointwise limit of the functions  $\phi_n$  as  $n \rightarrow +\infty$ . So to prove the subadditivity of  $\phi$ , it suffices to prove the subadditivity of all  $\phi_n$ , which we do by induction on  $n$ . The case  $n = 0$  is trivial, so we proceed to the induction step from  $n$  to  $n + 1$ . Let  $x, y \in \mathbb{R}$ ,  $x \geq y$ . Here we consider several cases.

*Case 1:*  $y \leq 0$ . This case is trivial as  $\phi_{n+1}$  is monotone.

*Case 2:*  $y \geq \frac{1}{3}$ . In this case,

$$\phi_{n+1}(x + y) \leq 1 = \frac{1}{2} + \frac{1}{2} \leq \phi_{n+1}(x) + \phi_{n+1}(y).$$

*Case 3:*  $x \leq \frac{1}{3}$ . As  $x, y$  and  $x + y$  are all  $\leq \frac{2}{3}$ , we have

$$\begin{aligned} \phi_{n+1}(x + y) &= \frac{1}{2} \cdot \phi_n(3x + 3y) \\ &\leq \frac{1}{2} \cdot \phi_n(3x) + \frac{1}{2} \cdot \phi_n(3y) = \phi_{n+1}(x) + \phi_{n+1}(y). \end{aligned}$$

---

Received by the editors September 2, 1994 and, in revised form, May 11, 1995.

1991 *Mathematics Subject Classification.* Primary 26D15.

*Key words and phrases.* Subadditive functions.

*Case 4:*  $0 \leq y \leq \frac{1}{3} \leq x$ . As  $x + y \geq \frac{1}{3}$ , we have

$$\begin{aligned}\phi_{n+1}(x+y) &= \frac{1}{2} + \frac{1}{2} \cdot \phi_n(3x+3y-2) \\ &\leq \frac{1}{2} + \frac{1}{2} \cdot \phi_n(3x-2) + \frac{1}{2} \cdot \phi_n(3y) = \phi_{n+1}(x) + \phi_{n+1}(y).\end{aligned}$$

These four cases exhaust all the possibilities, so the proof is complete.  $\square$

Let us recall that a modulus of continuity is a function  $f$  defined, continuous, nondecreasing and subadditive on  $[0, 1]$  with  $f(0) = 0$ .

**Corollary.** *The standard Cantor function is a modulus of continuity.*

The author wishes to thank the referee for his useful suggestions.

#### REFERENCES

1. Darst, R.B., *Some Cantor sets and Cantor functions*, Math. Mag. **45** (1972), 2–7. MR **46:311**
2. Matkowski, J. and Świątkowski, T., *On subadditive functions*, Proc. Amer. Math. Soc. **119** (1993), 187–197. MR **93k:26002**
3. Richey, M., *Mapping the Cantor set onto  $[0, 1]$  : a new construction*. preprint.
4. Rosenbaum, R.A., *Subadditive functions*, Duke Math. J. **17** (1950), 227–247. MR **12:164a**

DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY. 042 00 KOŠICE, SLOVAKIA  
E-mail address: `dobos@ccsun.tuke.sk`